

## 老年肌少症及其营养干预研究进展

郭宇, 赵前程, 李莹, 邱斌, 刘佳雯

### Research Progress on Senile Sarcopenia and Its Nutritional Intervention

GUO Yu, ZHAO Qiancheng, LI Ying, QIU Bin, and LIU Jiawen

在线阅读 View online: <https://doi.org/10.13386/j.issn1002-0306.2023070013>

## 您可能感兴趣的其他文章

### Articles you may be interested in

#### 河北省核桃坚果蛋白质、脂肪及脂肪酸组成分析

Analysis of Protein, Fat and Fatty Acid Composition of Walnut Resources in Hebei Province

食品工业科技. 2021, 42(13): 292-298 <https://doi.org/10.13386/j.issn1002-0306.2020070374>

#### 云南3种胶质食用菌营养成分分析与蛋白质营养价值评价

Analysis of Nutritional Components and Evaluation of Protein Nutritional Value of Three Kinds of Gelatinous Edible Fungi in Yunnan

食品工业科技. 2021, 42(14): 328-333 <https://doi.org/10.13386/j.issn1002-0306.2020090143>

#### 益生菌对糖尿病干预作用的研究进展

Research progress on the interventional effects of probiotics on diabetes

食品工业科技. 2017(22): 321-324 <https://doi.org/10.13386/j.issn1002-0306.2017.22.062>

#### 多酚与蛋白质相互作用研究方法进展

Progress in Research Methods for the Interaction between Polyphenols and Proteins

食品工业科技. 2018, 39(24): 340-345 <https://doi.org/10.13386/j.issn1002-0306.2018.24.058>

#### 蛋白质热聚集行为机理及其对蛋白质功能特性影响的研究进展

Research Progress on the Mechanism of Protein Thermal Aggregation Behavior and Its Influence on Functional Properties of Protein

食品工业科技. 2018, 39(24): 318-325 <https://doi.org/10.13386/j.issn1002-0306.2018.24.054>

#### 贵州不同品种芸豆营养品质差异分析

Analysis on the Difference of Nutritional Quality of Different Kidney Bean Varieties in Guizhou

食品工业科技. 2021, 42(18): 73-80 <https://doi.org/10.13386/j.issn1002-0306.2021010107>



关注微信公众号, 获得更多资讯信息

郭宇, 赵前程, 李莹, 等. 老年肌少症及其营养干预研究进展 [J]. 食品工业科技, 2024, 45(7): 370–377. doi: 10.13386/j.issn1002-0306.2023070013

GUO Yu, ZHAO Qiancheng, LI Ying, et al. Research Progress on Senile Sarcopenia and Its Nutritional Intervention[J]. Science and Technology of Food Industry, 2024, 45(7): 370–377. (in Chinese with English abstract). doi: 10.13386/j.issn1002-0306.2023070013

· 专题综述 ·

# 老年肌少症及其营养干预研究进展

郭宇<sup>1,2</sup>, 赵前程<sup>1,\*</sup>, 李莹<sup>1</sup>, 邱斌<sup>2,3,\*</sup>, 刘佳雯<sup>3</sup>

(1.大连海洋大学食品科学与工程学院, 辽宁大连 116023;

2.山东省农业科学院农产品加工与营养研究所, 山东济南 250000;

3.枣庄市东粮生物科技发展有限公司, 山东枣庄 277200)

**摘要:** 肌少症是一种与年龄相关的肌肉质量和功能下降疾病, 极大增加了老年人骨折、摔伤和死亡风险。其发病机制主要涉及运动神经元缺失、激素失衡和炎症因子与胰岛素抵抗等因素。补充蛋白质、氨基酸、n-3、n-6 多不饱和脂肪酸以及维生素等营养素可以有效防治肌少症。文章着重对肌少症的发病机制和营养干预进行了阐述, 期为老年肌少症研究提供参考。

**关键词:** 老年, 肌少症, 营养干预, 蛋白质, 脂肪酸

中图分类号: TS201.4

文献标识码: A

文章编号: 1002-0306(2024)07-0370-08

DOI: 10.13386/j.issn1002-0306.2023070013



本文网刊:

## Research Progress on Senile Sarcopenia and Its Nutritional Intervention

GUO Yu<sup>1,2</sup>, ZHAO Qiancheng<sup>1,\*</sup>, LI Ying<sup>1</sup>, QIU Bin<sup>2,3,\*</sup>, LIU Jiawen<sup>3</sup>

(1.School of Food Science and Engineering, Dalian Ocean University, Dalian 116023, China;

2.Institute of Food & Nutrition Science and Technology Shandong Academy of Agricultural Sciences,

Jinan 250000, China;

3.Zaozhuang Dongliang Biotechnology Development Co., Ltd., Zaozhuang 277200, China)

**Abstract:** Sarcopenia, an age-associated loss of muscle mass and function, greatly increases the risk of fractures, falls, and death in older adults. The pathogenesis of which is mainly involves motor neuron loss, hormone imbalance, inflammatory factors, and insulin resistance, etc. Supplementing nutrients such as protein, amino acid, n-3, n-6 polyunsaturated fatty acids and vitamin and other nutrients can effectively prevent and treat the sarcopenia. This paper focuses on the pathogenesis and nutritional intervention of sarcopenia to provide a reference for the research of sarcopenia.

**Key words:** old age; sarcopenia; nutritional intervention; protein; fatty acid

肌少症又称肌肉减少症, 是指与年龄相关的肌肉质量减少, 同时存在肌肉力量和/或躯体功能下降的老年综合征<sup>[1]</sup>。随着老龄化问题日益严重, 老年病患者越来越多, 该症作为发病率较高的骨骼肌运动系统疾病, 近年来已成为研究重点<sup>[2]</sup>。Liu 等<sup>[3]</sup>通过荟萃分析(Meta)明确了肌少症是社区老年人全因死亡

率的一个重要预测因子。然而现阶段我国对肌少症的了解还处于初步阶段, 对老年肌少症的风险认知不够深入, 防治不够规范<sup>[4]</sup>。提高民众对肌少症的认知率并对其进行有效防治, 是当下亟待解决的问题。

肌少症的发病机制与运动神经元丢失、胰岛素抵抗、炎症因子、线粒体功能障碍及营养不足等因素

收稿日期: 2023-07-04

基金项目: 山东省农业科学院农业科技创新工程; 营养功能型豆制品加工与副产物资源化利用 (CXGC2023A38); 2022 年度山东重点扶持区域引进急需紧缺人才项目“基于小麦精深加工关键技术的糖尿病患者专用食品研究开发”。

作者简介: 郭宇 (1999-), 女, 硕士研究生, 研究方向: 食品营养与安全, E-mail: 1026295879@qq.com。

\* 通信作者: 赵前程 (1966-), 男, 博士, 教授, 研究方向: 食品营养与安全, E-mail: qczhao@dlou.edu.cn。

邱斌 (1982-), 男, 博士, 副研究员, 研究方向: 营养感知与食品创制, E-mail: qiubin2009@live.cn。

有关<sup>[5-10]</sup>。其防治方式主要有运动干预、药物治疗和营养补充<sup>[11]</sup>。研究表明,基于阻力的运动对受试者的肌肉力量和功能有积极影响<sup>[12]</sup>。Hsu 等<sup>[13]</sup>的分析显示运动可以改善肌少症肥胖患者的肌肉力量和身体功能。老年患者肌肉力量不足,行动不便,难以有效完成运动训练,因此运动干预并非防治肌少症的最佳办法。药物治疗则应根据老年人多重用药安全管理专家共识和用药指南进行科学、规范的药品知识宣教,且多重用药会对健康造成严重影响<sup>[14]</sup>。而营养干预在提供充足能量的同时,又可以通过营养补充改善老年人肌肉质量,安全方便。因营养素缺乏是导致肌少症的重要原因之一,临床上的治疗方案有补充蛋白质、氨基酸和维生素 D,也有研究发现其他营养素如多不饱和脂肪酸<sup>[15]</sup>和矿物质<sup>[16]</sup>等与肌少症防治有着密切联系。近年来,肌少症研究现状的探讨或着重发病机制,或着重干预方式,本文整合了肌少症发病机制和营养干预的相关资料并进行探讨,以期分析其危害和营养因素来提高民众认知,并为肌少症的防治提供参考。

## 1 老年肌少症及其发病机制研究进展

### 1.1 老年肌少症

肌少症首次由 Rosenberg 于 1989 年命名。2010 年欧洲老年肌少症工作组首先发表了肌少症共识<sup>[17]</sup>,建议对于肌少症的诊断同时使用肌肉质量下降和肌肉功能或强度下降两个指标,并应用这些特征进一步将其概念定义为“骨骼肌前减少症”、“肌少症”和“严重肌少症”。2011 年国际肌少症工作组对肌少症进行了明确定义<sup>[1]</sup>。2014 年亚洲肌少症工作组在发表的共识报告中推荐了一种动态的肌少症研究方法,通过测量肌肉质量、力量和功能、身体表现、虚弱状态、日常生活活动、在给定时间内的日常生活基本活动、害怕跌倒以及尿失禁作为肌少症研究的结果指标,为亚洲人提供了肌少症的有效诊断<sup>[18]</sup>。2018 年 11 月国际肌少症与衰弱工作组发布了国际肌少症临床实践指南,内容包括肌少症的筛查、诊断及治疗,指出可侧重蛋白质与维生素 D 的研究,适度的蛋白质摄入可作为一种非药物干预的治疗方案<sup>[19]</sup>。2019 年中华医学会老年医学分会发布了老年人肌少症口服营养补充中国专家共识(2019)<sup>[20]</sup>,指出营养缺失的肌少症患者可进行口服营养补充。2021 年中华医学会老年医学分会、《中华老年医学杂志》编辑委员会编写了适合我国老年人肌少症患者的诊疗规范共识<sup>[4]</sup>。分别对肌少症定义、流行病学、诊断参数、诊断切点值、筛查和诊断流程以及预防和治疗发表了十五条推荐。

### 1.2 发病机制

肌少症是年龄增加相关性疾病,其危险因素与发病机制多种多样。源于组织学、生物化学和分子生物学的研究证实,神经元丢失、激素水平变化、线粒体功能障碍、营养素缺乏或摄入改变等均与肌少

症有关,这些均是衰老相关的多因素综合作用的结果。探明肌少症发生与发展的机理有助于其临床干预,但目前尚无明确的首要致病因素。

1.2.1 衰老与运动神经元丢失 随着年龄增长,老年人机体骨骼肌逐渐萎缩,患肌少症概率增大。老年人肌肉质量的减少,主要是由于运动神经元尤其是  $\alpha$  运动神经元( $\alpha$ -MNs)的逐渐丧失导致肌纤维数量下降。 $\alpha$ -MNs 丢失,神经肌肉连接(NMJs)改变,特别是 II 型肌纤维向 I 型肌纤维转变可导致肌纤维的萎缩,同时伴有肌内和肌间脂肪浸润导致的骨骼肌结构异常,最终发展为肌少症<sup>[5-6]</sup>。在骨骼肌再生过程中起主要作用的卫星细胞由于衰老而数量减少,导致肌肉的修复和再生能力降低<sup>[21]</sup>。还有研究表明衰老引起的线粒体损害可以加速活性氧和细胞能量缺乏,损伤的线粒体不断累积可触发运动神经元死亡<sup>[22]</sup>。

1.2.2 激素失衡 人体内多种激素可通过不同方式影响骨骼肌。研究表明,雌激素是一种抗氧化剂和肌膜稳定剂,可以积极影响骨骼肌的收缩特性并防止肌肉损伤。雌激素替代疗法可能间接影响骨骼肌蛋白质的更新,从而降低人体肌肉质量和力量损失<sup>[23]</sup>。Briocche 等<sup>[24]</sup>发现生长激素替代治疗可导致骨骼肌蛋白合成、线粒体生物发生途径的增加和骨骼肌再生、蛋白质降解抑制因子的降低来预防肌少症。另有研究比较了肌少症患者和非肌少症患者血液和激素指标,发现生长激素(growth hormone, GH)和胰岛素样生长因子(insulin-like growth factor, IGF-1)与老年肌少症有关,且 IGF-1 与骨骼肌质量的减少独立相关,IGF-1 表型是激活骨骼肌卫星细胞以启动骨骼肌细胞修复并促进其增殖的主要因素<sup>[25]</sup>。

1.2.3 胰岛素抵抗与炎症因子增多 胰岛素抵抗通过影响蛋白质合成途径以及激活蛋白质降解途径,最终导致肌肉损失<sup>[26]</sup>。Kim 等<sup>[7]</sup>通过人群实验得出胰岛素抵抗指数 HOMR-IR 与骨骼肌质量指数 SMI 呈负相关的结论。

炎症因子增多是推进肌少症发生发展的重要机制之一。随着年龄增大,炎症因子肿瘤坏死因子  $\alpha$  (tumor necrosis factor, TNF- $\alpha$ )、白细胞介素-6(interleukin-6, IL-6)和 C 反应蛋白(C-reactive protein, CRP)表达增加影响蛋白质的合成,促进蛋白质的分解,最终导致骨骼肌质量减少<sup>[27]</sup>。研究发现较高水平的循环炎症标志物(包括 CRP、TNF- $\alpha$  和 IL-6)与较低水平的骨骼肌力量及质量显著相关<sup>[8]</sup>。另有研究通过测定受试者的 IL-6、CRP、下肢肌肉质量和肌肉蛋白水解量,证实了炎症增加了骨骼肌蛋白水解,降低了骨骼质量和力量<sup>[28]</sup>。

1.2.4 线粒体功能障碍 肌细胞中线粒体完整性的丧失是导致肌肉退化的主要因素。线粒体功能的调节可诱导衰老细胞死亡,进而改善肌肉骨骼的健康、质量和功能。有证据表明,线粒体干预电子传递链的多个调控步骤,诱导了促炎衰老相关分泌表型元件的



表达,如 IL-6 和 IL-8,这些促炎细胞因子已被发现可以改变卫星细胞的基因表达程序,深刻影响肌肉再生,并导致肌肉功能的年龄依赖性下降<sup>[9]</sup>。还有研究表明,老年人线粒体膜通透性转换孔功能的失调导致线粒体活性氧(ROS)的过量产生、线粒体 DNA 释放增加参与肌肉萎缩的泛素连接酶的表达和促凋亡因子的释放等促进肌肉质量下降和功能丧失<sup>[29]</sup>。

1.2.5 营养素缺乏 Shan 等<sup>[10]</sup>在调研中发现营养素缺乏是肌少症发病的一个重要预测因子。蛋白质合成、降解和自噬参与肌少症的分子机制,老年人营养素摄入不足时,如蛋白质的缺乏使肌肉蛋白质的降解速率超过合成速率时,肌肉质量损失、肌肉功能逐渐减退<sup>[30]</sup>。维生素 D 缺乏也会因转录因子 FOXO1 持续激活,诱导胰岛素抵抗从而降低肌肉质量<sup>[31]</sup>。营养与老年人的肌肉质量、力量和功能有密切联系,这表明营养在肌少症的预防和管理方面发挥着重要作用,应充分补充蛋白质、维生素 D、抗氧化营养素和长链多不饱和脂肪酸以确保老年人的能量和营养需求<sup>[32]</sup>。

虽然老年肌少症发病机制的研究已经取得长足进步,但目前许多研究仍停留在探索阶段,新型特异性指标的发现,对诊断标准的制订及针对性治疗的开展都将具有重要意义,基因组学、蛋白组学、代谢组学等多组学的研究及大数据分析能为老年肌少症的发病机制研究及特异指标挖掘提供保障。

## 2 老年肌少症的营养干预

肌少症的病因与营养素的摄入不足密切相关。有效正确的营养补充对肌肉质量有着积极的影响,是预防肌少症最有效的方法。补充蛋白质、脂肪酸和多种维生素等营养素有益于肌肉组织合成,对抗肌少症效果显著。在难以通过膳食满足肌少症患者营养需求的情况下,额外的营养补充是必不可少的干预手段。

### 2.1 蛋白质对肌少症的干预

蛋白质代谢是营养问题的核心,蛋白质摄入不足也是肌少症患者营养不良的首要因素。研究表明,有效的蛋白摄入,可显著增加肌肉蛋白储备,减少肌肉量流失,从而有效控制老年肌少症病情发展<sup>[33]</sup>。Morifuji 等<sup>[34]</sup>研究发现与膳食酪蛋白相比,膳食乳清蛋白增加了肝脏葡萄糖激酶和骨骼肌糖原合酶的活性,研究也首次证明了基于乳清蛋白的饮食增加大鼠骨骼肌中的糖原含量,还观察到乳清蛋白通过不同的机制调节糖原代谢。最佳蛋白质摄入量与肌肉功能测定结果有关,如骨量、骨骼肌量、力量和患病风险<sup>[35]</sup>。欧洲肠外肠内营养学会推荐:对于身体状况良好的老人每日蛋白质适宜摄入量为 1.0~1.2 g/kg;急慢性老年患者 1.2~1.5 g/kg,其中优质蛋白质如动物蛋白和乳清蛋白的比例最好占一半<sup>[36]</sup>。Elisabet 等<sup>[37]</sup>通过实验得出,增加乳清蛋白的摄入量比单独的碳水化合物会更有效地促进肌肉蛋白的合成。还有文章指出,在老年肌少症的营养配方中乳清蛋白的

比例应占 60% 及以上<sup>[38]</sup>。

### 2.2 肽及氨基酸对肌少症的干预

肽是由两个或者两个以上氨基酸组成的化合物,作为蛋白质的水解中间产物和发挥活性作用的基团,肽与肌少症也有着密切联系。尤莉蓉<sup>[39]</sup>通过长达 6 个月的人群对照实验确定了大豆水解蛋白肽可以使人体肌肉含量显著增加。有研究通过动物实验得出低分子量的核桃蛋白水解肽具有显著的抗氧化(增强超氧化物歧化酶(SOD)、GSH-Px 和过氧化氢酶(CAT)活性,降低丙二醛(MDA)含量)和免疫调节(提高免疫器官指数,促进肠道 T 淋巴细胞增殖和 sIgA 分泌)等预防肌肉衰减的功能<sup>[40]</sup>。汪云凤等<sup>[41]</sup>也通过动物实验证实玉米低聚肽通过降低肌环指蛋白-1 和肌萎缩 F 盒蛋白的 mRNA 表达量,抑制肌肉萎缩。目前,德国公司嘉利达开发了一款可帮助肌少症患者改善肌肉状况的胶原蛋白肽(BODYBALANCE<sup>®</sup>),且经实验证明,产品中的蛋白肽可使受试者的肌力增加<sup>[42]</sup>。

氨基酸是蛋白质营养与代谢的基本单位。补充部分必需氨基酸、支链氨基酸或氨基酸代谢产物均有助于肌少症患者身体机能的改善。亮氨酸、缬氨酸和异亮氨酸统称为支链氨基酸(BCAA),支链氨基酸通过促进胰岛素和生长激素的释放来加快合成代谢和肌肉增长。Hernández 等<sup>[43]</sup>对 32 名经诊断的肝硬化和肌少症患者进行了为期 12 周的安慰剂和 BCAA 营养干预,结果显示 BCAA 组的锌水平高于安慰剂组,白蛋白水平也显著提高。因此补充 BCAA 可改善肝硬化伴随肌少症患者的肌肉质量。Park 等<sup>[44]</sup>对 54 名患者进行营养干预实验,单因素分析结果显示,受试者在补充 BCAA 和强化康复治疗后有更高的骨骼肌指数。Izumi 等<sup>[45]</sup>对肌少症老年人进行为期 8 周的 BCAA 干预,结果显示 BCAA 可以改善患者的肌肉状态。王蕾等<sup>[46]</sup>的动物实验表明亮氨酸和谷氨酰胺的联合可以延缓肌少症。我国青岛一项 1140 受试者参与的调查研究结果表明:较高的 BCAA 摄入量与较低的肌少症风险有关,这可能有益于预防肌少症并改善老年人的身体机能<sup>[47]</sup>。此外,其他氨基酸也对肌少症有着积极影响。补充瓜氨酸能增加氧化能量周转,降低 pH 功率比和 ATP 成本,改善骨骼肌代谢和/或收缩效率<sup>[48]</sup>。研究发现,天冬氨酸可以抑制脂多糖诱导的肌肉萎缩相关基因的表达,从而抑制蛋白质降解,改善肌肉状态<sup>[49]</sup>。还有动物实验表明,赖氨酸的补充也可以通过抑制蛋白水解提高骨骼肌质量<sup>[50]</sup>。

亮氨酸改善肌肉质量和力量的关键之一就是转化为代谢产物  $\beta$ -羟基- $\beta$ -甲基丁酸( $\beta$ -hydroxy- $\beta$ -methylbutyrate, HMB)。研究<sup>[51]</sup>表明, HMB 诱导急性肌肉合成代谢,增加肌蛋白的合成。Ellis 等<sup>[52]</sup>研究表明, HMB 可通过刺激蛋白质合成和减少蛋白质降解,同时减少炎症,从而改善肌肉损失。还有文章

通过归纳分析法汇总了 HMB 可以通过调节靶蛋白的表达水平、激活丝氨酸-苏氨酸激酶(MAPK)和磷脂酰肌醇 3-激酶(PI3K)的信号通路、影响半胱天冬酶和 ROS 的产生以及转化为 HMG-CoA 来抑制骨骼肌蛋白质的降解<sup>[53]</sup>。美国雅培开发了一款类似的乳清蛋白固体饮料——安补素, 其中添加了两大有助肌肉健康的成分 CaHMB 和乳清蛋白。

与年轻人相比, 老年人的代谢效率下降, 因此需要更多的蛋白质摄入, 以保证蛋白质合成。作为综合治疗的一部分, 补充足量的优质蛋白质或肽有助于预防甚至逆转肌肉减少症。同时饮食中增加富含亮氨酸的食材或额外补充特殊氨基酸同样对肌少症有较好的干预效果。

### 2.3 脂肪酸对肌少症的干预

脂肪酸是人类生命活动中的重要物质, 不同种类脂肪酸均有其特定功能和用途。多不饱和脂肪酸与人体健康密不可分, 在人体生理中起着极为重要的代谢作用, 根据其结构分为 n-6 和 n-3 两大主要系列。n-3 多不饱和脂肪酸、n-6 多不饱和脂肪酸和共轭亚油酸(CLA)在肌少症的形成中发挥着不同作用。n-3 多不饱和脂肪酸和 CLA 主要通过降低炎症反应、机体氧化活性, 影响睾酮生物合成、胰岛素敏感性和前列腺素产量和直接参与肌肉蛋白合成与降解来延缓肌少症的形成, 同时还可通过增加抗阻力运动的效果及与蛋白质、肌酸等营养物质的联合作用来延缓肌少症的形成。n-6 多不饱和脂肪酸主要是促使肌肉蛋白分解, 降低炎症反应来抑制肌少症的形成<sup>[15,54-60]</sup>。Calder 等<sup>[15]</sup>指出 n-3 和 n-6 多不饱和脂肪酸的抗炎机制可以抑制促炎转录因子的激活, 并提到 n-3 脂肪酸补充剂可降低内毒素刺激的单核细胞或单核细胞对 TNF、IL-1 $\beta$  和 IL-6 的产生, 而炎症可能是骨骼肌蛋白减少和消耗的重要机制。Rahman 等<sup>[55]</sup>研究结果显示 CLA 可以通过维持衰老过程中的氧化还原平衡来预防肌少症。Cintia 等<sup>[61]</sup>研究表明富含 n-3 多不饱和脂肪酸的补充剂组能有效提高老年妇女的肌肉激活水平并改善肌肉力量和功能。Mizanur 等<sup>[59]</sup>发现添加 CLA 可以通过调节炎症和破骨细胞因子的标志物来防止骨骼和肌肉质量的损失。Virtanen 等<sup>[62]</sup>研究发现 n-6 多不饱和脂肪酸与关键炎症标志物 CRP 呈强负相关。此外, 中链脂肪酸甘油三酯(MCT)对骨骼肌也有着积极影响。MCT 通常是由 6~12 个碳原子的饱和脂肪酸包括己酸(C6:0)、庚酸(C7:0)、辛酸(C8:0)、壬酸(C9:0)、癸酸(C10:0)和月桂酸(C12:0)组成的甘油三酯<sup>[63]</sup>。Sakiko 等<sup>[64]</sup>研究表明, 晚餐时联合补充 6 g 剂量的 MCT, 可改善虚弱老年人的肌肉力量和功能, 且通过补充 MCT, 脂肪酸被大量用于骨骼肌的能量生产。MCT 供能可达到葡萄糖的两倍多, 还可增强人体对钙、镁和氨基酸的吸收, 从而促进氨基酸和矿物质对肌少症患者肌肉质量的改善。

目前, 脂肪酸与肌少症的关系研究主要集中于多不饱和脂肪酸, 但大多是流行病学资料、动物实验以及多不饱和脂肪酸与其他营养物质的联合作用的人群干预实验, 单独分析多不饱和脂肪酸作用的人群干预实验报道较少。单类脂肪酸或不同比例脂肪酸复配与老年肌少症在临床应用的组效、量效、构效关系及其作用机制仍需要更进一步的研究。

### 2.4 维生素与微量元素对肌少症的干预

维生素可通过不同机制参与肌少症的预防和治疗。维生素 D 是目前具备最多理论研究和临床实践支撑的能有效防治肌少症的维生素。Girgis 等<sup>[65]</sup>发现维持正常的维生素 D 信号传递是防止肌肉功能和肌肉大小丧失的有效策略。维生素 D 也可通过基因组和非基因组机制精细控制骨骼肌细胞<sup>[66]</sup>。25(OH)D3 是维生素 D 的重要中间代谢产物, 也是维生素 D 在人体内储存的重要形式。相关研究表明, 其水平降低将导致肌肉质量下降, 进而增加肌少症患病风险<sup>[67]</sup>。《老年人肌少症口服营养补充中国专家共识(2019)》推荐老年肌少症患者维生素 D 的补充剂量为 15~20  $\mu\text{g}/\text{d}$ (600~800 U/d)<sup>[20]</sup>。维生素 E 可以通过增强抗氧化能力、抑制炎症反应、减轻氧化磷脂和稳定胰岛素结构等来增加损伤骨骼肌的存活率和增强肌肉再生能力<sup>[68]</sup>。研究表明维生素 E 的膳食摄入量与骨骼肌质量呈正相关<sup>[69]</sup>。Bo 等<sup>[70]</sup>还发现补充维生素 E 可以有效改善老年肌少症患者的肌肉质量和力量。一项针对 55~56 岁的成年人观察研究结果揭示在女性人群中, 较低的维生素 K 水平与较弱的握力、小腿围及躯体功能相关<sup>[71]</sup>。但维生素 K 在骨骼肌中的具体作用机制目前还有待进一步研究。维生素 B<sub>12</sub> 在外周神经和中枢神经的整合过程中起着重要的作用。Bulut 等<sup>[72]</sup>通过观察记录 403 位患者的行走速度、肌力和肌肉质量, 分别对每个患者进行 4 米行走实验、握力实验和生物阻抗实验, 并进行维生素 B<sub>12</sub> 检测, 结果揭示肌少症可能与维生素 B<sub>12</sub> 缺乏症有关。维生素 C 具有很强的还原性, 是有效的抗氧化剂, 也是生物合成和基因调节酶家族的辅助因子。研究发现饮食中的维生素 C 与中、老年人骨骼肌质量呈正相关, 表明膳食维生素 C 摄入可能有助于减少与年龄相关的肌肉损失<sup>[73]</sup>。Welch 等<sup>[69]</sup>研究表明, 维生素 C 可以降低衰老过程中骨骼肌质量和力量的损失, 进而防治肌少症。

血清低水平微量元素是老年衰弱的独立危险因素, 并且衰弱风险会随微量元素缺乏而增大。一项研究探讨了老年人血清硒与骨骼肌质量的关系, 发现通过调整年龄、性别、生活方式、身体和代谢因素, 较低血清硒水平与低骨骼肌质量的高风险相关<sup>[74]</sup>。另有研究对 331 名受试者的数据分析结果显示, 与非肌少症受试者相比, 肌少症受试者消耗的许多微量营养素(钾、镁、铁、钙等)显著减少, 表明不平衡的饮食可能与肌少症和肌肉骨骼健康不良有关, 但还需要前



瞻性研究来确认这些因素<sup>[16]</sup>。

维生素广泛参与细胞增殖、骨骼肌蛋白质合成、氧化损伤及修复等过程,与人体肌肉健康密切相关,但其对肌肉减少症的相关作用机制尚未完全明确。今后需进一步深入探讨维生素及微量元素影响肌肉减少症发生发展的机制。

### 3 总结与展望

肌少症主要引起老年人运动功能障碍,生活质量下降,增加家庭及社会的经济负担。因其发病可能性强、人群广,目前已经被社会逐渐重视,治疗这种疾病对提高老年人生活质量有很重要的意义。老年肌少症的发病机制涉及衰老、激素和营养缺失等多个方面。充足的优质蛋白、维生素D和脂肪酸等营养干预可以提前预防或及时改善肌肉减少症。

从肌肉减少症概念的提出开始,国际上对此综合征的研究已有二十多年,在定义、发病机制、诊断和营养干预措施方面都有一定的积累,但还是存在许多空白亟需相关学科专业或产业的重视和投入。一是发病机制的研究仍应进一步深入。通过基因组学、蛋白组学、代谢组学等多组学的研究及大数据分析,挖掘新型特异性指标,对诊断标准的制订及针对性治疗的开展都将具有重要意义。二是营养干预手段应进一步完善。营养干预等是对肌少症干预的重要手段,其相关研究将是今后预防和治疗肌少症的重要研究方向,后续仍需拓展并完善单一营养素或复合营养素对老年肌少症的发病机制及防治作用研究。三是针对肌少症的多元化营养健康食品产业需要大力发展。随着经济发展和健康型消费模式的出现,营养健康食品的产业规模年均增幅也达到了10%~15%,已成为未来食品行业的重要发展方向。针对肌肉减少症人群的特殊营养需求,开展多营养素互作、营养素随加工工艺的动态变化规律的研究,最终开发出适合肌少症人群专用的特殊医学用途配方食品、营养强化食品或功能性食品等多元化产品,大力促进我国营养健康产业发展,助力国民健康。

© The Author(s) 2024. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

### 参考文献

[1] ROGER A F, BRUNO V, WILLIAM J E, et al. Sarcopenia: An undiagnosed condition in older adults. Current consensus definition: Prevalence, etiology, and consequences. International working group on sarcopenia[J]. *Journal of the American Medical Directors Association*, 2011, 12(4): 249–256.

[2] CRUZ-JENTOFT A J, SAYER A A. Sarcopenia[J]. *The Lancet*, 2019, 393(10191): 2636–2646.

[3] LIU Ping, QIU Kuihao, SHAN Hai, et al. Sarcopenia as a predictor of all-cause mortality among community-dwelling older people: A systematic review and meta-analysis[J]. *Maturitas*, 2017, 103: 16–22.

[4] 刘娟,丁清清,周白,等.中国老年人肌少症诊疗专家共识(2021)[J]. *中华老年医学杂志*, 2021, 40(8): 943–952. [LIU Juan, DING Qingqing, ZHOU Bai, et al. Expert consensus on the diagnosis and treatment of sarcopenia in the elderly in China (2021)[J]. *Chinese Journal of Geriatrics*, 2021, 40(8): 943–952.]

[5] JIA Yuyin, QIAN Zhiyuan, CHEN Yuqi, et al. MicroRNA regulatory networks in the pathogenesis of sarcopenia[J]. *Journal of Cellular and Molecular Medicine*, 2020, 24(9): 4900–4912.

[6] LARS L, DEGENS H, LI M, et al. Sarcopenia: Aging-related loss of muscle mass and function[J]. *Physiological Reviews*, 2019, 99(1): 427–511.

[7] KIM T N, PARK M S, LIM K I, et al. Relationships between sarcopenic obesity and insulin resistance, inflammation, and vitamin D status: The Korean sarcopenic obesity study[J]. *Clinical Endocrinology*, 2013, 78(4): 525–532.

[8] CAMILLA S L T, LACHLAN A N T, ANDREA B M. Markers of inflammation and their association with muscle strength and mass: A systematic review and meta-analysis[J]. *Ageing Research Reviews*, 2020, 64: 101185.

[9] FERRI E, EMANUELE M, RICCARDO C, et al. Role of age-related mitochondrial dysfunction in sarcopenia[J]. *International Journal of Molecular Sciences*, 2020, 21(15): 5236.

[10] SHAN Hai, CAO Li, WANG Hui, et al. Association between sarcopenia and nutritional status and physical activity among community-dwelling Chinese adults aged 60 years and older[J]. *Geriatrics & Gerontology International*, 2017, 17(11): 1959–1966.

[11] 李悦梵,董娟,张任飞,等.肌少症的治疗进展[J]. *中国临床保健杂志*, 2023, 26(3): 313–317. [LI Yuefang, DONG Juan, ZHANG Renfei, et al. Progress in the treatment of sarcopenia[J]. *Chinese Journal of Clinical Care*, 2023, 26(3): 313–317.]

[12] FELICITY R W, BERZIGOTTI A, LORD J M, et al. Review article: Impact of exercise on physical frailty in patients with chronic liver disease[J]. *Alimentary Pharmacology & Therapeutics*, 2019, 50(9): 988–1000.

[13] HSU K J, LIAO C D, TSAI M W, et al. Effects of exercise and nutritional intervention on body composition, metabolic health, and physical performance in adults with sarcopenic obesity: A meta-analysis[J]. *Nutrients*, 2019, 11(9): 2163.

[14] 张文婧,王佳贺.老年肌少症与营养干预的研究进展[J]. *国际老年医学杂志*, 2020, 41(2): 125–128. [ZHANG Wenjing, WANG Jiahe. Research progress on sarcopenia and nutritional intervention in the elderly[J]. *International Journal of Geriatrics*, 2020, 41(2): 125–128.]

[15] CALDER P C. Marine omega-3 fatty acids and inflammatory processes: Effects, mechanisms and clinical relevance[J]. *BBA-Molecular and Cell Biology of Lipids*, 2015, 1851(4): 469–484.

[16] CHARLOTTE B, LOCQUET M, TOUVIER M, et al. Association between dietary nutrient intake and sarcopenia in the SarcoPhAge study[J]. *Ageing Clinical and Experimental Research*, 2019, 31(6): 815–824.

[17] CRUZ-JENTOFT A J, BAEYENS J P, BAUER J M, et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European working group on sarcopenia in older people[J]. *Age and Ageing*, 2010, 39(4): 412–423.

[18] LIANG Kungchen, LIU Likuo, WOO J, et al. Sarcopenia in Asia: Consensus report of the Asian working group for sarcopenia [J]. *Journal of the American Medical Directors Association*, 2014, 15(2): 95–101.

[19] DENT E, MORLEY J E, CRUZ-JENTOFT A J, et al. Inter-

- national clinical practice guidelines for sarcopenia (ICFSR): Screening, diagnosis and management[J]. *The Journal of Nutrition, Health & Aging*, 2018, 22(10): 1148–1161.
- [20] 中华老年医学杂志编辑委员会, 中华医学会老年医学分会, 老年人肌少症口服营养补充中国专家共识(2019)[J]. *中华老年医学杂志*, 2019, 38(11): 1193–1197. [Chinese Journal of Geriatrics, Editorial Board. Geriatrics Branch of Chinese Medical Association. Chinese expert consensus on oral nutritional supplementation for sarcopenia in the elderly (2019)[J]. *Chinese Journal of Geriatrics*, 2019, 38(11): 1193–1197.]
- [21] HIROKI N, SHINYA F, AKIR A A, et al. Pathophysiology and mechanisms of primary sarcopenia (Review)[J]. *International Journal of Molecular Medicine*, 2021, 48(2): 1–8.
- [22] SONJAK V, JACOB K J, SPENDIFF S, et al. Reduced mitochondrial content, elevated reactive oxygen species, and modulation by denervation in skeletal muscle of prefrail or frail elderly women [J]. *The Journals of Gerontology: Series A*, 2019, 74(12): 1887–1895.
- [23] HANSEN M. Female hormones: Do they influence muscle and tendon protein metabolism?[J]. *Proceedings of the Nutrition Society*, 2017, 77(1): 32–41.
- [24] BRIOCHE T, KIREEV R A, CUESTA S, et al. Growth hormone replacement therapy prevents sarcopenia by a dual mechanism: Improvement of protein balance and of antioxidant defenses [J]. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 2014, 69(10): 1186–1198.
- [25] AILIN B, MA Y, ZHOU X Z, et al. Association between sarcopenia and levels of growth hormone and insulin-like growth factor-1 in the elderly[J]. *BMC Musculoskeletal Disorders*, 2020, 21(1): 1–9.
- [26] RASTOGI K R, MARK C, LUIGI F. Age-related and disease-related muscle loss: The effect of diabetes, obesity, and other diseases[J]. *The Lancet Diabetes & Endocrinology*, 2014, 2(10): 819–829.
- [27] POWERS S K, SMUDER A J, JUDGE A R. Oxidative stress and disuse muscle atrophy: Cause or consequence?[J]. *Current Opinion in Clinical Nutrition and Metabolic Care*, 2012, 15(3): 240.
- [28] MASAYUKI K, HARADA M, NOYAMA S, et al. Association between inflammation and skeletal muscle proteolysis, skeletal mass and strength in elderly heart failure patients and their prognostic implications[J]. *BMC Cardiovascular Disorders*, 2020, 20(1): 1–9.
- [29] PHILIPPE J, LEDUC G, HUSSAIN S N A, et al. Mitochondrial dynamics and mitophagy in skeletal muscle health and aging [J]. *International Journal of Molecular Sciences*, 2021, 22(15): 8179.
- [30] RONG Shuang, WANG Liangliang, PENG Zhao, et al. The mechanisms and treatments for sarcopenia: Could exosomes be a perspective research strategy in the future?[J]. *Journal of Cachexia, Sarcopenia and Muscle*, 2020, 11(2): 348–365.
- [31] CHEN Songcang, VILLALTA S A, AGRAWAL D K. FOXO1 mediates vitamin D deficiency-induced insulin resistance in skeletal muscle[J]. *Journal of Bone and Mineral Research: The Official Journal of the American Society for Bone and Mineral Research*, 2016, 31(3): 585–595.
- [32] ROBINSON M S, REGINSTER J Y, RIZZOLI R, et al. Does nutrition play a role in the prevention and management of sarcopenia?[J]. *Clinical Nutrition*, 2018, 37(4): 1121–1132.
- [33] 张林英, 梅群超, 张炜琦, 等. 抗阻训练联合高乳蛋白营养对老年肌肉减少症患者肌张力及生活质量的影响[J]. *国际老年医学杂志*, 2023, 44(2): 196–199. [ZHANG Linying, MEI Qunchao, ZHANG Weiqi, et al. Effects of resistance training combined with high milk protein nutrition on muscle tone and quality of life in elderly patients with sarcopenia[J]. *International Journal of Geriatrics*, 2023, 44(2): 196–199.]
- [34] MORIFUJI M, KENSUKE S, CHIAKI S, et al. Dietary whey protein increases liver and skeletal muscle glycogen levels in exercise-trained rats[J]. *British Journal of Nutrition*, 2005, 93(4): 439–445.
- [35] AGNES N P, TOMMY C. Health effects of protein intake in healthy elderly populations: A systematic literature review[J]. *Food & Nutrition Research*, 2014, 58(1): 23364.
- [36] TIELAND M, MARLOU L D, NIKITA V D Z, et al. Protein supplementation increases muscle mass gain during prolonged resistance-type exercise training in frail elderly people: A randomized, double-blind, placebo-controlled trial[J]. *Journal of the American Medical Directors Association*, 2012, 13(8): 713–719.
- [37] ELISABET B, AARSLAND A, WOLFE R R. Effect of an amino acid, protein, and carbohydrate mixture on net muscle protein balance after resistance exercise[J]. *International Journal of Sport Nutrition and Exercise Metabolism*, 2004, 14(3): 255–271.
- [38] YVETTE C L, EVAN A, THOMAS L, et al. Protein type and caloric density of protein supplements modulate postprandial amino acid profile through changes in gastrointestinal behaviour: A randomized trial[J]. *Clinical Nutrition*, 2016, 35(1): 48–58.
- [39] 尤莉蓉. 大豆肽运动补剂的研发及其促肌肉增长作用分析[J]. *食品研究与开发*, 2017, 38(8): 163–165. [YOU Lirong. Research and development of soybean peptide sports supplement and its role in promoting muscle growth[J]. *Food Research and Development*, 2017, 38(8): 163–165.]
- [40] LI Fang, REN Dayong, CUI Lingyu, et al. Antifatigue, antioxidant and immunoregulatory effects of peptides hydrolyzed from manchurian walnut (*Juglans mandshurica* Maxim.) on mice[J]. *Grain & Oil Science and Technology*, 2018, 1(1): 44–52.
- [41] 汪云凤, 闫九明, 周正明, 等. 玉米低聚肽改善地塞米松诱导的肌肉萎缩效果研究[J]. *华西医学*, 2023, 38(3): 429–437. [WANG Yunfeng, YAN Jiuming, ZHOU Zhengming, et al. Effect of zee oligopeptide on improving dexamethasone-induced muscle atrophy[J]. *West China Medical Journal*, 2023, 38(3): 429–437.]
- [42] ZDZIEBLIK D, STEFFEN O, MANFRED W B, et al. Collagen peptide supplementation in combination with resistance training improves body composition and increases muscle strength in elderly sarcopenic men: A randomised controlled trial[J]. *British Journal of Nutrition*, 2015, 114(8): 1237–1245.
- [43] HERNÁNDEZ C M, LLOP E, GÓMEZ P L, et al. Adding branched-chain amino acids to an enhanced standard-of-care treatment improves muscle mass of cirrhotic patients with sarcopenia: A placebo-controlled trial[J]. *American Journal of Gastroenterology*, 2021, 116(11): 2241–2249.
- [44] PARK M K, LEE S J, CHOI E, et al. The effect of branched chain amino acid supplementation on stroke-related sarcopenia[J]. *Frontiers in Neurology*, 2022, 13: 744945.
- [45] IZUMI T, YOSHIMURA Y, SHIMAZU S, et al. Effects of branched-chain amino acids and vitamin D supplementation on physical function, muscle mass and strength, and nutritional status in sarcopenic older adults undergoing hospital-based rehabilitation: A multicenter randomized controlled trial[J]. *Geriatrics & Gerontology International*, 2019, 19(1): 12–17.

- [46] 王蕾, 陈强谱, 蔡婷婷, 等. 亮氨酸+谷氨酰胺、中药对梗阻性黄疸大鼠肌少症的干预研究[C]//中国营养学会, 中国疾病预防控制中心营养与健康所, 农业农村部食物与营养发展研究所, 中国科学院上海营养与健康研究所, 华中科技大学公共卫生学院. 中国营养学会第十五届全国营养科学大会论文汇编. 2022: 789. [WANG Lei, CHEN Qiangpu, CAI Tingting, et al. Intervention of leucine+glutamine and traditional Chinese medicine on sarcopenia in rats with obstructive jaundice[C]//Chinese Nutrition Society, Institute of Nutrition and Health, Chinese Center for Disease Control and Prevention, Institute of Food and Nutrition Development, Ministry of Agriculture and Rural Affairs, Shanghai Institute of Nutrition and Health, Chinese Academy of Sciences, School of Public Health, Huazhong University of Science and Technology. Proceedings of the 15th National Conference of Nutrition Science of Chinese Nutrition Society. 2022: 789.]
- [47] MA Wenjun, ZHANG Wei, ZHAI Jie, et al. Branched chain and other amino acid intakes are inversely associated with sarcopenia among community elders in Qingdao, China[J]. *Asia Pacific Journal of Clinical Nutrition*, 2022, 31(1): 97–107.
- [48] ALBERTO C G, PASCUAL F J, NORIEGA G D C, et al. L-citrulline supplementation and exercise in the management of sarcopenia[J]. *Nutrients*, 2021, 13(9): 3133.
- [49] LIU Yulan, WANG Xiuying, LENG Weibo, et al. Aspartate inhibits LPS-induced MAFbx and MuRF1 expression in skeletal muscle in weaned pigs by regulating Akt, AMPK $\alpha$  and FOXO1[J]. *Innate Immunity*, 2017, 23(1): 34–43.
- [50] TOMONORI S, ITO Y, NAGASAWA T. Dietary l-lysine suppresses autophagic proteolysis and stimulates Akt/mTOR signaling in the skeletal muscle of rats fed a low-protein diet[J]. *Journal of Agricultural and Food Chemistry*, 2015, 63(37): 8192–8198.
- [51] STOUT J R, SMITH-RYAN A E, FUKUDA D H, et al. Effect of calcium  $\beta$ -hydroxy- $\beta$ -methylbutyrate (CaHMB) with and without resistance training in men and women 65+ yrs: A randomized, double-blind pilot trial[J]. *Experimental Gerontology*, 2013, 48(11): 1303–1310.
- [52] ELLIS A C, HUNTER G R, GOSS A M, et al. Oral supplementation with beta-hydroxy-beta-methylbutyrate, arginine, and glutamine improves lean body mass in healthy older adults[J]. *Journal of Dietary Supplements*, 2019, 16(3): 281–293.
- [53] 田浩楠, 汪军, 张锋.  $\beta$ -羟基- $\beta$ -甲基丁酸盐治疗老年肌少症的分子机制研究进展[C]//中国营养学会, 中国疾病预防控制中心营养与健康所, 农业农村部食物与营养发展研究所, 中国科学院上海营养与健康研究所, 华中科技大学公共卫生学院. 中国营养学会第十五届全国营养科学大会论文汇编. 2022: 573. [TIAN Haonan, WANG Jun, ZHANG Feng. Research progress on the molecular mechanism of  $\beta$ -hydroxy- $\beta$ -methylbutyrate in the treatment of sarcopenia in the elderly[C]//Chinese Nutrition Society, Institute of Nutrition and Health, Chinese Center for Disease Control and Prevention, Institute of Food and Nutrition Development, Ministry of Agriculture and Rural Affairs, Shanghai Institute of Nutrition and Health, Chinese Academy of Sciences, School of Public Health, Huazhong University of Science and Technology. Proceedings of the 15th National Conference of Nutrition Science of Chinese Nutrition Society. 2022: 573.]
- [54] FILIPPO M, BARONE R, CATANESE P, et al. Do fat supplements increase physical performance?[J]. *Nutrients*, 2013, 5(2): 509–524.
- [55] RAHMAN M, HALADE G V, EL J A, et al. Conjugated linoleic acid (CLA) prevents age-associated skeletal muscle loss[J]. *Biochemical and Biophysical Research Communications*, 2009, 383(4): 513–518.
- [56] MONTAÑO M, ZIMMER K E, DAHL E, et al. Effects of mixtures of persistent organic pollutants (POPs) derived from cod liver oil on H295R steroidogenesis[J]. *Food and Chemical Toxicology*, 2011, 49(9): 2328–2335.
- [57] NALL J L, WU G Y, KIM K H, et al. Dietary supplementation of L-arginine and conjugated linoleic acid reduces retroperitoneal fat mass and increases lean body mass in rats[J]. *The Journal of Nutrition*, 2009, 139(7): 1279–1285.
- [58] RAHMAN M M, HALADE G V, WILLIAMS P J, et al. t10c12-CLA maintains higher bone mineral density during aging by modulating osteoclastogenesis and bone marrow adiposity[J]. *Journal of Cellular Physiology*, 2011, 226(9): 2406–2414.
- [59] MIZANUR R M, ARUNABH B, JAMEELA B, et al. Conjugated linoleic acid protects against age-associated bone loss in C57BL/6 female mice[J]. *The Journal of Nutritional Biochemistry*, 2006, 18(7): 467–474.
- [60] SMITH G I, ATHERTON P, REEDS D N, et al. Dietary omega-3 fatty acid supplementation increases the rate of muscle protein synthesis in older adults: A randomized controlled trial[J]. *The American Journal of Clinical Nutrition*, 2011, 93(2): 402–412.
- [61] CINTIA L N R, RODACKI A L F, PEREIRA G, et al. Fish-oil supplementation enhances the effects of strength training in elderly women[J]. *The American Journal of Clinical Nutrition*, 2012, 95(2): 428–436.
- [62] VIRTANEN J K, MURSU J, VOUTILAINEN S, et al. The associations of serum n-6 polyunsaturated fatty acids with serum C-reactive protein in men: The kuopio ischaemic heart disease risk factor study[J]. *European Journal of Clinical Nutrition*, 2018, 72(3): 342–348.
- [63] RAY A M, LARRY THOMPSON D V M, GEORGE A B. Lack of toxicity by medium chain triglycerides (MCT) in canines during a 90-day feeding study[J]. *Food and Chemical Toxicology*, 2008, 47(1): 35–39.
- [64] SAKIKO A, EZAKI O, SUZUKI M. Medium-chain triglycerides in combination with leucine and vitamin D increase muscle strength and function in frail elderly adults in a randomized controlled trial[J]. *The Journal of Nutrition*, 2016, 146(5): 1017–1026.
- [65] GIRGIS C M, CHA K M, SO B, et al. Mice with myocyte deletion of vitamin D receptor have sarcopenia and impaired muscle function[J]. *Journal of Cachexia, Sarcopenia and Muscle*, 2019, 10(6): 1228–1240.
- [66] VASANT H, CUMMING R G, NAGANATHAN V, et al. Longitudinal associations between vitamin D metabolites and sarcopenia in older Australian men: The concord health and aging in men project[J]. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 2017, 73(1): 131–138.
- [67] 王静, 赵红敏, 张宇, 等. 血清 25-羟维生素 D<sub>3</sub> 水平与 2 型糖尿病肾病患者并发肌少症的相关性分析[J]. *中国医刊*, 2023, 58(3): 333–336. [WANG Jing, ZHAO Hongmin, ZHANG Yu, et al. Correlation between serum 25-hydroxy vitamin D<sub>3</sub> level and myalopenia in patients with type 2 diabetic nephropathy[J]. *Chinese Medical Journal*, 2023, 58(3): 333–336.]
- [68] CHUNG E, MO H, WANG S. Potential roles of vitamin E in age-related changes in skeletal muscle health[J]. *Nutrition Research*, 2018(49): 23–36.
- [69] WELCH A A, JENNINGS A, KELAIDITI E, et al. Cross-sectional associations between dietary antioxidant vitamins C, E and



carotenoid intakes and sarcopenic indices in women aged 18-79 years[J]. *Calcified Tissue International*, 2020, 106(4): 331-342.

[ 70 ] BO Yacong, LIU Changfeng, JI Zhe. A high whey protein, vitamin D and E supplement preserves muscle mass, strength, and quality of life in sarcopenic older adults: A double-blind randomized controlled trial[J]. *Clinical Nutrition*, 2019, 38(1): 159-164.

[ 71 ] NANA N, KOMATSU J, IWASE H, et al. Effects of the combination of vitamin K and teriparatide on the bone metabolism in ovariectomized rats[J]. *Biomedical Reports*, 2015, 3(3): 295-300.

[ 72 ] BULUT E A, SOYSAL P, AYDIN A E, et al. Vitamin B12

deficiency might be related to sarcopenia in older adults[J]. *Experimental Gerontology*, 2017, 95: 136-140.

[ 73 ] LEWIS L N, HAYHOE R P G, MULLIGAN A A, et al. Lower dietary and circulating vitamin C in middle- and older-aged men and women are associated with lower estimated skeletal muscle mass[J]. *The Journal of Nutrition*, 2020, 150(10): 2789-2798.

[ 74 ] CHEN Y L, YANG K C, CHANG H H, et al. Low serum selenium level is associated with low muscle mass in the community-dwelling elderly[J]. *Journal of the American Medical Directors Association*, 2014, 15(11): 807-811.